



Development and Validation of the Short-Form Ryff's Psychological Well-Being Scale for Clinical Nurses in Taiwan

Tony Szu-Hsien Lee¹, Hueyi-Fang Sun², Hui-Hsun Chiang²

¹Department of Health Promotion and Health Education, National Taiwan Normal University, ²School of Nursing, National Defense Medical Center, Taipei, Taiwan

Background: Research into clinical nurses' psychological well-being (PWB) is limited and fragmented. A reliable and valid measure is needed to advance the field of nurses' mental health. **Objective:** To examine the psychometric validity of Short-Form PWB Scale (PWBSs) among clinical nurses. **Methods:** This study adopted a cross-sectional design. Randomized cluster sampling was used to recruit clinical nurses from a medical center in Taipei, Taiwan, from July to October 2015. Nurses were recruited if they worked for more than 3 months in the medical center. The 84-item PWBS was used to assess PWB. Based on exploratory and confirmatory factor analysis in this study, an 18-item version of Ryff's PWBS was developed using 474 respondents. The Short-Form PWBS was tested for internal consistency, construct validity, and criterion-related validity. **Results:** Cronbach's alpha was 0.88, with aggregated subscale alphas of 0.72–0.88, except 0.57 for autonomy. Good construct validity and criterion-related validity of the Short-Form PWBS were found. **Discussion:** The study results showed that the Short-Form PWBS is suitable and recommended for Taiwanese clinical nurses.

Key words: Clinical nurses, construct validity, criterion-related validity, internal consistency, psychological well-being

INTRODUCTION

Nurses' psychological well-being (PWB) is positively associated with quality of care and job satisfaction.¹ A systematic review shows that poor well-being is associated with moderate-to-high levels of burnout, and poor well-being among health-care staff contributes to poorer quality of care and less patient safety.² The original version of Ryff's PWB Scale (PWBS) consists of six dimensions, namely autonomy, environmental mastery, purpose in life, positive relationships, personal growth, and self-acceptance. The scale has 192 items totally, and each dimension has 32 items.³

Later, different versions of the PWBS (3 items, 7 items, 9 items, and 14 items per subscale) were developed because there was a need for a short and easy-to-take version since clinical staff such as physicians and nurses have limited time to complete items.⁴ Based on the literature review, most studies focused on the dimensions of depression and emotional exhaustion. Few addressed nurses' positive emotions such as PWB.⁵ Hence, the study aimed to examine the psychometric integrity of a short form of the Mandarin PWBS in a sample of clinical nurses.

Received: January 01, 2019; Revised: January 10, 2019;
Accepted: January 21, 2019

Corresponding Author: Dr. Hui-Hsun Chiang, No. 161, Sec. 6, Minquan E RD, Neihu dist. Taipei City, 11490, Taiwan.
Tel: +886-2-87923100#18761; Fax: 886-2-66005702.

E-mail: sheisvivan@gmail.com

METHODS

Ethical considerations

This study protocol was approved by the Institutional Review Board of the Tri-Service General Hospital (Approval No. 1-105-05-109).

Design

The study adopted a cross-sectional design to examine the validity and reliability of the Chinese version of PWBS in clinical nurses in Taiwan.

Participants and procedure

A cluster random sampling with probability proportionate to sample size was used to recruit clinical nurses from a medical center in Taipei, Taiwan, from July 2015 to October 2015. We stratified the sample by specialty to obtain similar number of participants at each work unit. Details of the study procedure

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

How to cite this article: Lee TS, Sun HF, Chiang HH. Development and validation of the Short-Form Ryff's Psychological Well-Being Scale for clinical nurses in Taiwan. J Med Sci 2019;XX:XX-XX.

are described in a previous article.⁶ Specifically, the inclusion criterion was working at least 3 months in a clinical setting. The exclusion criteria were nonshift working, a part-time job, or a nonclinical staff position.

A sample of clinical nurses ($n = 474$) agreed to participate in the study after the study purpose and procedure were explained by the researchers. Participants were assured that their participation was anonymous and would not influence their work. After signing the consent form, it took about 20–30 min for the participants to complete the questionnaire in a quiet room with comfortable air condition and temperature. Two separate boxes were used to collect the consent forms and questionnaires to assure anonymity.

Instruments

Demographic items

Demographic items include gender, age, marital status, education level, work experience, perceived health, and unit.

Ryff's Psychological Well-Being Scale

The PWBS has 84 items, 14 items in each of the six subscales, which included autonomy, environmental mastery, personal growth, positive relations with others, purpose in life, and self-acceptance.⁴ The Mandarin version was confirmed to have good validity and reliability ($\alpha = 0.89$) among adults.⁷ Internal consistency was 0.88 for the short version in this study. Respondents answer each item using a 6-point scale, ranging from 1 ("strongly disagree") to 6 ("strongly agree")."

Statistical analyses

Descriptive data were analyzed using **SPSS** software version 20.0. Scores for the PWBS are presented as means, standard deviations (SDs), Cronbach's alpha, Pearson's correlation coefficients, and F values from analysis of variance.

To determine the psychometric validity of the Short-Form PWBS, exploratory factor analysis (EFA) and confirmatory factor analysis (CFA) were performed using **Mplus software version 5.1** and a criterion significance level of $P < 0.05$.⁸ EFA was performed to reduce the items to a common factor pattern. The uni-dimensionality of the resulting scales was examined through CFA. Finally, the reliability of these scales was assessed through internal consistency analysis.⁹ Items were selected on the basis of their fit with the factor pattern (percentage of total variance and factor loadings) and conformance to the guiding theoretical definition of PWB dimensions. One factor per dimension is recommended in the literature, and results from the scree plot confirmed that the number of factors is appropriate. The indicators of model fit used for this study included: (1) comparative fit index (CFI) ≥ 0.90 ,¹⁰ (2) root mean square error of approximation (RMSEA) < 0.07 ,¹¹ and (3) standardized root mean squared residual (SRMR) ≤ 0.08 .¹⁰

RESULTS

Sample characteristics

The participants' background characteristics are listed in Table 1. Most participants were female ($n = 455$, 96%), and the mean age was 29.6 years ($SD = 7.4$). Approximately 77% were single or divorced ($n = 363$) and 96.4% had an educational level of bachelor's degree or higher ($n = 447$). About one-third reported that their work experience was < 2 years and 61.2% ($n = 280$) had been employed at the hospital for < 5 years. Most participants worked in medical-surgical wards ($n = 282$, 59.2%) and 46.3% reported no disturbed health ($n = 220$). The sum scores for the dimensions ranged from 3 to 18. There was no years-of-experience difference for gender, education level, and PWBS scores, except for the autonomy subscale [Table 1].

PSYCHOMETRIC VALIDITY OF THE 18-ITEM PSYCHOLOGICAL WELL-BEING SCALE

Principal axis factoring with varimax rotation was performed to examine item loadings. The Kaiser–Meyer–Olkin measures of sampling adequacy were all > 0.76 , indicating that there was a sufficient level of factorability. In addition, Bartlett's test of sphericity was statistically significant at $P < 0.001$, indicating that the correlation matrix was not identical to the factor structure matrix. Overall, the data were appropriate for factor analysis. An initial examination of the scree plot suggested six factors. Eigenvalues of the selected items were > 1 and communality values were > 0.5 , except for autonomy subscale (about 0.3).¹²

Results from the EFA, including components matrix, communality values, total variance explained, and rule of cross-loading, collectively suggested that three items (short version) comprising a one-factor solution were most appropriate. Internal consistency (alpha coefficients) for the reduced scales ranged from 0.72 to 0.81, except for autonomy subscale (0.57), and correlations of the subscales with the 84-item total scale ranged from 0.70 to 0.87. Table 2 presents the EFA results for the three items in each PWBS subscale.

Then, CFA was performed to examine the uni-dimensionality of the PWBS. The second-order factor analysis model-fit statistics for the originally proposed model are as follows: CFI = 0.78, RMSEA = 0.11, and SRMR = 0.26, indicating that the six-factor model did not fit very well. After determining the relation between positive relations with others and self-acceptance, the revised model-fit statistics were found to be as follows: CFI = 0.90, RMSEA = 0.076, and SRMR = 0.048. Standardized factor loadings for the PWBS dimensions ranged

Table 1: Sample characteristics of clinical nurses (n=474)

| Characteristics | n (%) | | | P |
|--------------------------------|------------------|---------------------|-------------|--------|
| | Total (n=474) | Years of experience | | |
| | | <2 years | ≥2 years | |
| Age (years), mean (SD) | 29.6 (7.4) | 24.4 (2.71) | 32.2 (7.4) | |
| Gender | | | | |
| Female | 455 (96.0) | 155 (32.8) | 300 (63.3) | 0.48 |
| Male | 19 (4.0) | 5 (1.0) | 14 (3.0) | |
| Education level | | | | |
| Senior high school | 2 (0.4) | 0 | 2 (0.4) | 0.05 |
| College or university | 457 (96.4) | 159 (33.5) | 298 (62.9) | |
| Graduate school | 15 (3.2) | 1 (0.2) | 14 (3.0) | |
| Workplace | | | | |
| ER | 54 (11.4) | 14 (3.0) | 40 (8.4) | <0.01 |
| OR | 30 (6.3) | 9 (1.9) | 21 (4.4) | |
| ICU | 108 (22.8) | 25 (5.3) | 83 (17.5) | |
| Medical-surgical ward | 282 (59.2) | 112 (23.6) | 170 (35.9) | |
| Marital status | | | | |
| Married | 111 (23.4) | 3 (.6) | 108 (22.8) | <0.001 |
| Single/divorced/separated | 363 (76.6) | 157 (33.1) | 206 (43.5) | |
| Years of experience | | | | |
| <2 | 160 (33.8) | | | |
| 2-5 | 130 (27.4) | | | |
| 5-10 | 84 (17.7) | | | |
| 11-15 | 43 (9.1) | | | |
| >5 | 57 (12.1) | | | |
| Perceived health | | | | |
| Very disturbed (often) | 14 (3.0) | 5 (1.1) | 9 (1.9) | 0.01 |
| Disturbed (usually) | 178 (37.6) | 42 (8.9) | 136 (28.7) | |
| Not disturbed (sometimes) | 220 (46.3) | 93 (19.6) | 127 (26.7) | |
| Good (not at all) | 62 (13.1) | 20 (4.2) | 42 (8.9) | |
| PWBS | 72.9 (9.6) | 72.43 (9.1) | 73.19 (9.8) | 0.41 |
| Autonomy | 11.3 (1.9) | 10.9 (1.9) | 11.5 (1.9) | <0.01 |
| Environment mastery | 12.5 (2.0) | 12.5 (2.0) | 12.5 (2.0) | 0.70 |
| Personal growth | 13.3 (1.9) | 13.1 (2.0) | 13.3 (1.9) | 0.18 |
| Positive relations with others | 12.3 (2.8) | 12.2 (3.0) | 12.3 (2.6) | 0.81 |
| Purpose in life | 12.2 (2.2) | 12.4 (2.1) | 12.1 (2.2) | 0.13 |
| Self-acceptance | 11.4 (2.6) | 11.3 (2.7) | 11.5 (2.5) | 0.39 |

ER=Emergency room; OR=Operation room; ICU=Intensive care unit; PWBS=Psychological Well-Being Scale; SD=Standard deviation

from 0.84 to 0.99, except for the error term for the relation between positive relations with Others (loading = 0.42) and self-acceptance (loading = 0.43), as shown in Table 2. Correlation coefficients for relationships between the 18-item and 84-item PWBS and the corresponding subscales ranged from 0.70 to 0.87. Cronbach's alpha coefficients for the

six subscales of the 18-item PWBS ranged from 0.72 to 0.81, except 0.57 for autonomy subscale. The overall alpha coefficient for the 18-item PWBS was 0.88. There was a positive correlation between perceived health and the 18-item PWBS ($r = 0.20$, $P < 0.001$), indicating that criterion validity is acceptable.

AQ14 Table 2: Summary of dimensions and reliability of the Short-Form Psychological Well-Being Scale ($n=474$)

| Subscale and item number | f^a | Percent total variance (%) | h^2 | f^b | r | Cronbach's alpha |
|--------------------------------|-------|----------------------------|-------|-------|-------|------------------|
| PWBS | | | | | | 0.88 |
| Autonomy | | 44.03 | | 0.84* | 0.70* | 0.57 |
| 8 | 0.51 | 7.73 | 0.26 | 0.43* | | |
| 14 | 0.53 | 20.82 | 0.28 | 0.46* | | |
| 50 | 0.52 | 15.49 | 0.27 | 0.72* | | |
| Environmental mastery | | 55.01 | | 0.99* | 0.76* | 0.72 |
| 51 | 0.62 | 7.06 | 0.39 | 0.66* | | |
| 69 | 0.69 | 32.53 | 0.45 | 0.71* | | |
| 81 | 0.62 | 15.43 | 0.39 | 0.69* | | |
| Personal growth | | 54.35 | | 0.87* | 0.71* | 0.81 |
| 46 | 0.71 | 8.08 | 0.50 | 0.73* | | |
| 52 | 0.72 | 31.05 | 0.51 | 0.83* | | |
| 70 | 0.71 | 15.21 | 0.50 | 0.73* | | |
| Positive relations with others | | 57.49 | | 0.42* | 0.87* | 0.75 |
| 13 | 0.69 | 36.08 | 0.47 | 0.66* | | |
| 31 | 0.65 | 7.66 | 0.42 | 0.73* | | |
| 61 | 0.68 | 13.75 | 0.46 | 0.73* | | |
| Purpose in life | | 55.63 | | 0.96* | 0.74* | 0.77 |
| 23 | 0.68 | 6.54 | 0.46 | 0.67* | | |
| 47 | 0.62 | 16.34 | 0.47 | 0.74* | | |
| 53 | 0.63 | 32.74 | 0.47 | 0.77* | | |
| Self-acceptance | | 55.63 | | 0.43* | 0.80* | 0.75 |
| 42 | 0.60 | 8.48 | 0.36 | 0.77* | | |
| 66 | 0.63 | 31.09 | 0.40 | 0.71* | | |
| 84 | 0.62 | 14.15 | 0.38 | 0.64* | | |

AQ13 * $P < 0.001$. ^aExploratory factor analysis with principal axis factoring was used to deduct items from each subscale, ^bConfirmatory factor analysis with principal axis factoring was used to verify the loadings of each dimension. f^a =Standardized item loading; f^b =Standardized factor loading; h^2 =Communality; r =Correlation with 84-item scale; PWBS=Psychological Well-Being Scale

DISCUSSION

Results from the study showed that the Short-Form (18 items) Mandarin PWBS has good psychometric integrity and thus is suitable for assessing the PWB of clinical nurses in Taiwan. The finding of moderate-to-strong associations of positive relations with others and self-acceptance is noteworthy. **The present study results are different from previous study results.**^{3,4,13} Self-acceptance is highly correlated with environment mastery, but only moderately correlated with positive relations with others. Nurses working in clinical practice feel stressful and exhausting. The relationships between nurses and the rest of the medical team are critical factors for job outcomes.¹⁴ By developing positive relations with others, in which empathy and intimacy are expressed, the nursing staff becomes more compassionate and understanding in their own lives and in their views of others.¹⁵ More self-compassion, consequently, is

associated with more compassion for others and the ability to understand that life is a journey, full of ups and downs, and not perfect. This, in turn, predicts a higher quality of professional life, which results in a higher level of self-acceptance.¹⁶

Our findings concerning the positive association of perceived health with PWB are consistent with previous results.^{17,18} A cross-sectional online survey of 697 nurses in Taipei showed that nurses with self-rated poor health were more likely to suffer from psychological symptoms than those in good health.¹⁷ Another survey of 432 volunteer nurses in South Korea revealed that increasing resources not only promoted healthy behavior but also improved stress management. The resulting higher levels of health responsibility and spiritual growth were found to improve nursing performance and care quality.¹⁹ Thus, perceived health in nurses is important not only for their own mental health, but also for the quality of patient care.

The subscales autonomy, environmental mastery, personal growth, purpose in life, positive relations with others, and self-acceptance performed well on all criteria and can be used independently. Autonomy performed well on construct validity but not so well on internal consistency. The finding that nurses with more than 2 years of experience showed higher levels of autonomy is consistent with a previous study.²⁰ Results from a sample of 3032 noninstitutionalized women and men between the ages of 25 and 74 years who were interviewed over telephone or completed a mail questionnaire showed that well-educated women with multiple roles had greater autonomy.²⁰ In the present study, most married nurses with more than 2 years of experience were not only capable of being a clinical care staff member, an educator, and a leader in the workplace, but also a wife, a daughter, and/or a mother **within the family**.

This study presents the advantages and limitations of quantitative analyses that should be taken into account for improving the clinical implications of PWB measurement. Both EFA and CFA were used to understand the dimensions of Short-Form PWBS. Moreover, a cluster random sampling with probability proportionate to sample size was used; the data were shown as descriptive terms and can be applied to a large number of participants. Several limitations should be taken into account, though the study is still noteworthy. The participants were mainly young women from a medical center, which limits the generalizability of the findings; these data do not reflect the national population. Further studies in multiple areas of the country are strongly recommended. Moreover, gender difference on the total PWBS and each subscale of PWBS should be assessed in the future.

IMPLICATIONS AND CONCLUSION

The 18-item short form of the Mandarin PWBS is an instrument with good reliability and validity that is suitable to measure the level of PWB in clinical nurses in Taiwan. The results contribute to the research literature on PWB among clinical nurses. This investigation can lead to the development of mental health promotion practices aligned with personal and institutional needs. The system may thus support PWB, which in turn, may result in a better nursing practice environment and higher quality of nursing care.

Acknowledgment

This research was supported by the Ministry of Science and Technology, R.O.C. (Grant No. 103-2314-B-016-042-). The authors would like to thank Professor Hillaire J Thompson for the English editing.

Financial support and sponsorship

This research was supported by the Ministry of Science and Technology, R.O.C. (Grant No. 103-2314-B-016-042-).

Conflicts of interest

There are no conflicts of interest.

REFERENCES

- Pahlevan Sharif S, Ahadzadeh AS, Sharif Nia H. Mediating role of psychological well-being in the relationship between organizational support and nurses' outcomes: A cross-sectional study. *J Adv Nurs* 2018;74:887-99.
- Hall LH, Johnson J, Watt I, Tsipa A, O'Connor DB. Healthcare staff wellbeing, burnout, and patient safety: A systematic review. *PLoS One* 2016;11:e0159015.
- Ryff CD. Happiness is everything, or is it? Explorations on the meaning of psychological well-being. *J Pers Soc Psychol* 1989;57:1069-81.
- Ryff CD, Keyes CL. The structure of psychological well-being revisited. *J Pers Soc Psychol* 1995;69:719-27.
- Arafa MA, Nazel MW, Ibrahim NK, Attia A. Predictors of psychological well-being of nurses in Alexandria, Egypt. *Int J Nurs Pract* 2003;9:313-20.
- Lee TS, Tzeng WC, Chiang HH. Impact of coping strategies on nurses' well-being and practice. *J Nurs Scholarsh* 2018. **In press.**
- Chiang HH, Chien LH, Lin JS, Yeh YH, Lee TS. Modeling psychological well-being and family relationships among retired older people in Taiwan. *Int J Ment Health Nurs* 2013;22:93-101.
- Muthén LK, Muthén BO. *Mplus User's Guide*. 5th ed. Los Angeles, CA: Muthén and Muthén; 1998-2007.
- Anderson JC, Gerbing DD. Structural equation modeling in practice: A review and recommended two-step approach. *Psychol Bull* 1988;103:411-23.
- Hu LL, Bentler PM. Cutoff criteria for fit indexes in covariance structure analysis: Conventional criteria versus new alternatives. *Struct Equ Modeling* 1999;6:1-55.
- Steiger JH. Understanding the limitations of global fit assessment in structural equation modeling. *Pers Individ Dif* 2007;42:893-8.
- Costello AB, Osborne JW. Best practices in exploratory factor analysis: Four recommendations for getting the most from your analysis. *Pract Assess Res Educ* 2005;10:1-9.
- Ryff CD. Psychological well-being in adult life. *Curr Dir Psychol Sci* 1995;4:99-104.
- Dos Santos Alves DF, da Silva D, de Brito Guirardello E. Nursing practice environment, job outcomes and safety

| | | |
|----|---|----|
| 1 | climate: A structural equation modelling analysis. J Nurs | 1 |
| 2 | Manag 2017;25:46-55. | 2 |
| 3 | 15. Westerhof GJ, Keyes CL. Mental illness and mental | 3 |
| 4 | health: The two continua model across the lifespan. | 4 |
| 5 | J Adult Dev 2010;17:110-9. | 5 |
| 6 | 16. Duarte J, Pinto-Gouveia J, Cruz B. Relationships between | 6 |
| 7 | nurses' empathy, self-compassion and dimensions of | 7 |
| 8 | professional quality of life: A cross-sectional study. Int J | 8 |
| 9 | Nurs Stud 2016;60:1-11. | 9 |
| 10 | 17. Chen MJ, Weng SS. Psychological symptoms among | 10 |
| 11 | hospital nurses in Taiwan: A cross-sectional study. BMC | 11 |
| 12 | | 12 |
| 13 | | 13 |
| 14 | | 14 |
| 15 | | 15 |
| 16 | | 16 |
| 17 | | 17 |
| 18 | | 18 |
| 19 | Author Queries??? | 19 |
| 20 | AQ1: Kindly provide revise CRF with adding journal title of all authors. | 20 |
| 21 | AQ2: Kindly check the typeset of the affiliation. | 21 |
| 22 | AQ3: Based of publication principles we cannot add or delete any author name once the article got accepted. | 22 |
| 23 | AQ4: Kindly check the E-mail id. | 23 |
| 24 | AQ6: Kindly provide missing abstract heading level of conclusion. | 24 |
| 25 | AQ7: Please provide complete manufacturer details such as company name, city, state and country. | 25 |
| 26 | AQ8: Please check the edit. | 26 |
| 27 | AQ9: Please check the insertion of text. | 27 |
| 28 | AQ10: Kindly check the heading. | 28 |
| 29 | AQ11: Kindly check the acknowledgement text. | 29 |
| 30 | AQ12: Please update reference, if possible, by providing complete publication details such as volume, year of publication, and page | 30 |
| 31 | range. | 31 |
| 32 | AQ13: Kindly check the redundant significant a and b. | 32 |
| 33 | AQ14: Kindly check the table formation | 33 |
| 34 | | 34 |
| 35 | | 35 |
| 36 | | 36 |
| 37 | | 37 |
| 38 | | 38 |
| 39 | | 39 |
| 40 | | 40 |
| 41 | | 41 |
| 42 | | 42 |
| 43 | | 43 |
| 44 | | 44 |
| 45 | | 45 |
| 46 | | 46 |
| 47 | | 47 |
| 48 | | 48 |
| 49 | | 49 |
| 50 | | 50 |
| 51 | | 51 |
| 52 | | 52 |